


# Exemplos | Examples

- Exemplo 1 | Example #1

Paciente masculino, 19 anos, com queixa de dor torácica. Sem comorbidades. Ao ECG, apresenta ritmo sinusal regular, QRS predominantemente negativo em DI e positivo em aVF, com eixo de de QRS entre  $+90^\circ$  e  $+180^\circ$ ; compatível com **desvio do eixo de QRS para direita**.


*19-year-old male patient presenting with chest pain. On ECG, tracing shows regular sinus rhythm with predominantly negative QRS in lead I and a positive QRS in lead aVF, with QRS axis between  $+90^\circ$  and  $+180^\circ$ ; consistent with **rightward deviation of the QRS axis**.*



- Exemplo 2 | Example #2

Paciente feminina, 25 anos, em avaliação pré-anestésica. Tabagista. Ao ECG, apresenta ritmo sinusal regular, complexos QRS predominantemente negativos em DI e positivo em aVF, com eixo de QRS entre  $+90^\circ$  e  $+180^\circ$ ; compatível com **desvio do eixo de QRS para direita**.

*25-year-old female patient undergoing a pre-anesthetic evaluation. Smoker. On ECG, tracing shows regular sinus rhythm with predominantly negative QRS complexes in lead I and positive in lead aVF and QRS axis between  $+90^\circ$  and  $+180^\circ$ ; corresponding to **rightward deviation of the QRS axis**.*



---

Revision #2

Created 22 May 2025 19:17:28 by Gabriela Miana

Updated 13 October 2025 12:57:30 by Gabriela Miana