Exemplos | Examples

- Exemplo 1 | Example #1

Paciente masculino, 68 anos, com queixa de dispneia. Tabagista, tem diagnósticos de diabetes mellitus e hipertensão arterial sistêmica. Em uso de diuréticos, hipoglicemiante oral, insulina e sinvastatina. Ao ECG, apresenta ritmo sinusal, extrassístole ventricular isolada e **intervalo QT prolongado**, representado por intervalo QT corrigido de 474 ms (>450 ms para homens), aumentado. Observa-se, também, área eletricamente inativa em parede em parede septal e alterações inespecíficas da repolarização ventricular.

*A 68-year-old male patient presenting with dyspnea. Smoker, with diagnosis of diabetes mellitus and hypertension. Currently taking diuretics, oral hypoglycemic agents, insulin, and simvastatin. On ECG, tracing shows sinus rhythm, isolated ventricular extrasystole and a **prolonged QT interval**, represented by a corrected QT interval of 474 ms (>450 for men). Electrically inactive area in the septal wall, and nonspecific ventricular repolarization abnormalities are also observed.*

AD_4nXcXD02THCPXfL7ZUeS0HvJvjFk_OXlC5T9P2DtpQc3ReMZzIGh4uUdNROBJDL1EnCfn8W4DVXUF

Revision #7
Created 2025-05-22 19:46:05 UTC by Gabriela Miana
Updated 2025-10-12 21:54:27 UTC by Gabriela Miana