


# Exemplos | Examples

## Exemplo 1 | Example #1

Paciente masculino, 91 anos, hipertenso controlado. Em uso de betabloqueador. Ao ECG, apresenta ausência de onda P e complexo QRS em um intervalo superior a 1,5s em DII longo, compatível com **pausa sinusal**. Observa-se, também, presença de BAV de 1º grau (PR > 200 ms em DII).


*91-year-old male patient with controlled hypertension on beta-blocker therapy. On ECG, tracing shows absence of P wave and QRS complex for an interval longer than 1.5 seconds in lead DII, consistent with **sinus pause**. First-degree heart block is also observed (PR interval > 200 milliseconds in lead II).*



- Exemplo 2 | Example #2

Paciente feminino, 68 anos, pós-PCR. Ao ECG, apresenta ausência de onda P e complexos QRS com duração de 140 ms (ritmo idioventricular acelerado) e presença de um intervalo superior a 1,5s em DII longo, correspondendo a **pausa sinusal**.

*68-year-old female patient, post-cardiopulmonary arrest. On ECG, tracing shows absence of P waves and QRS complexes with duration of 140 ms (accelerated idioventricular rhythm) with an interval greater than 1.5 seconds in long lead DII, corresponding to **sinus pause**.*



---

Revision #3

Created 22 May 2025 18:12:43 by Gabriela Miana

Updated 12 October 2025 21:53:22 by Gabriela Miana