


Exemplos | Examples

- Exemplo 1 | Example #1

Paciente do sexo masculino, 71 anos, com queixa de dor torácica e histórico de HAS, em uso de IECA ou BRA. Ao ECG, apresenta ritmo sinusal com episódio de **taquicardia atrial não sustentada (3 batimentos)** observado na derivação DII longo.


*Male patient, 71 years old, presenting with chest pain. Has been diagnosed with hypertension and is currently taking ACE inhibitors. On ECG, he presents sinus rhythm with an episode of **non-sustained atrial tachycardia** (3 beats) observed in the long DII lead.*



- Exemplo 2 | Example #2

Paciente feminina, 73 anos, com relato de dor torácica e histórico de HAS, em uso de atenolol e de maxopram. Ao ECG, apresenta bradicardia sinusal com frequência cardíaca de 48 bpm com extrassístolia supraventricular conduzida com QRS idêntico ao sinusal, isolada com episódio de **taquicardia atrial não sustentada**.

*Female patient, 73 years old, reports chest pain and history of hypertension, using atenolol and maxopram. On the ECG, sinus bradycardia with a heart rate of 48 bpm with supraventricular extrasystole conducted with a QRS identical to the sinus, isolated with an **episode of nonsustained atrial tachycardia**.*



Revision #3

Created 23 May 2025 14:04:03 by Gabriela Miana

Updated 13 October 2025 18:49:43 by Gabriela Miana